

# The Quadratic Formula

$$ax^2 + bx + c = 0$$

David

The L<sup>A</sup>T<sub>E</sub>X Project

2025-12-06

## Completing the Square

First, complete the square

- $ax^2 + bx + c = 0$

# Completing the Square

First, complete the square

- $ax^2 + bx + c = 0$

- $x^2 + \frac{b}{a}x + \frac{c}{a} = 0$

# Completing the Square

First, complete the square

- $ax^2 + bx + c = 0$

- $x^2 + \frac{b}{a}x + \frac{c}{a} = 0$

- $x^2 + \frac{b}{a}x + \left(\frac{b}{2a}\right)^2 = -\frac{c}{a} + \left(\frac{b}{2a}\right)^2$

# Completing the Square

First, complete the square

- $ax^2 + bx + c = 0$

- $x^2 + \frac{b}{a}x + \frac{c}{a} = 0$

- $x^2 + \frac{b}{a}x + \left(\frac{b}{2a}\right)^2 = -\frac{c}{a} + \left(\frac{b}{2a}\right)^2$

- $\left(x + \frac{b}{2a}\right)^2 = -\frac{c}{a} + \left(\frac{b}{2a}\right)^2$

## Solve for $x$

Now, solve for  $x$

- $\left(x + \frac{b}{2a}\right)^2 = -\frac{c}{a} + \left(\frac{b}{2a}\right)^2$

## Solve for $x$

Now, solve for  $x$

- $\left(x + \frac{b}{2a}\right)^2 = -\frac{c}{a} + \left(\frac{b}{2a}\right)^2$
- $x + \left(\frac{b}{2a}\right) = \pm \sqrt{-\frac{c}{a} + \left(\frac{b}{2a}\right)^2}$

## Solve for $x$

Now, solve for  $x$

- $\left(x + \frac{b}{2a}\right)^2 = -\frac{c}{a} + \left(\frac{b}{2a}\right)^2$
- $x + \left(\frac{b}{2a}\right) = \pm \sqrt{-\frac{c}{a} + \left(\frac{b}{2a}\right)^2}$
- $x = -\left(\frac{b}{2a}\right) \pm \sqrt{-\frac{c}{a} + \left(\frac{b}{2a}\right)^2}$

## Solve for $x$

Now, solve for  $x$

- $\left(x + \frac{b}{2a}\right)^2 = -\frac{c}{a} + \left(\frac{b}{2a}\right)^2$
- $x + \left(\frac{b}{2a}\right) = \pm \sqrt{-\frac{c}{a} + \left(\frac{b}{2a}\right)^2}$
- $x = -\left(\frac{b}{2a}\right) \pm \sqrt{-\frac{c}{a} + \left(\frac{b}{2a}\right)^2}$
- $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

# The Quadratic Formula

The solution of the quadratic equation.

$$ax^2 + bx + c = 0$$

where  $a$  is non-zero, is given by the formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$